

Simplified Methods for Construction, Assessment and Rapid Screening of Peptide Libraries in Bacteriophage

Rosemarie B. Christian†, Ronald N. Zuckermann, Janice M. Kerr
Liping Wang and Bruce A. Malcolm‡

Chiron Corporation
4560 Horton Street, Emeryville, CA 94608, U.S.A.

(Received 5 March 1992; accepted 2 June 1992)

An efficient strategy has been devised for the construction of diverse peptide libraries in bacteriophage vectors. This strategy was used to generate a library of 4×10^8 random decapeptide inserts in the pIII protein of bacteriophage fd. A novel method for evaluating the genetic diversity of bacteriophage libraries based on colony hybridization with partially degenerate oligonucleotides has been developed. The decapeptide library was affinity-selected with a previously characterized monoclonal antibody specific for the V3 loop of the gp120 protein of HIV-1. Immunological screening, an efficient technique for the rapid identification of putative binding bacteriophage, is described. Hexapeptide sequences similar to those obtained from affinity selection of a hexapeptide bacteriophage library were obtained from the decapeptide library in all five frames. Immunological screening of 20,000 clones from the two libraries after two rounds of affinity selection rapidly identified antibody-binding sequences; 93% and 86% of the sequences obtained from the hexapeptide and decapeptide libraries, respectively, had IC₅₀ values $\leq 10 \mu\text{M}$ as free peptides.

Keywords: bacteriophage; genetic diversity; immunological screening; peptide library

1. Introduction

The construction of diverse recombinant peptide libraries coupled with affinity selection and screening methods provide a powerful tool for the rapid identification of peptide ligands. Filamentous bacteriophage peptide libraries have been used to identify antibody epitopes (Scott & Smith, 1990; Cwirla *et al.*, 1990), and streptavidin-binding peptides (Devlin *et al.*, 1990). Phagemid libraries have been used to identify higher affinity variants of human growth hormone (Lowman *et al.*, 1991) as well as antibody epitopes (Felici *et al.*, 1991). Bacteriophage libraries displaying Fab fragments have been used to present the human (Marks *et al.*, 1991) and murine (Clackson *et al.*, 1991) immunological repertoires. Here, we describe an efficient strategy utilizing self-priming partially degenerate oligonucleotides for the generation of a bacteriophage peptide library of great size and complexity that should be applicable to the construction of any diverse library.

A colony hybridization technique was developed

to evaluate the genetic diversity of a decapeptide bacteriophage library. The genetic diversity of bacteriophage libraries has been examined by two different techniques to date. Cwirla *et al.* (1990) examined the diversity of an N-terminal hexapeptide library in pIII by analyzing the distribution of bases at each position within the codons *via* mass sequencing of pooled clones. Marks *et al.* (1991) have estimated the diversity of a V-gene library by PCR§ amplification of library colonies (Güssow & Clackson, 1989) followed by restriction analysis and sequencing. Both of these techniques yield a general qualitative picture of the genetic diversity of a phage library. The modified colony hybridization technique described here more rigorously assesses library diversity.

An immunological colony screening technique was utilized for the rapid screening of affinity-selected peptide libraries. Immunological screening has been used in the screening of cDNA libraries

† Present address: Microgenics Corporation, 2380A Bisso Lane, Concord, CA 94598

‡ Author to whom all correspondence should be addressed at: Department of Biochemistry, University of Alberta, Edmonton, Alberta T6G 2H7, Canada.

§ Abbreviations used: PCR, polymerase chain reaction; bp, base-pair(s); PEG, polyethylene glycol; u.v., ultraviolet light; TMAC, tetramethylammonium chloride; mAb, monoclonal antibody; PBS, phosphate-buffered saline; Ig, immunoglobulin; IC₅₀, concentration required for 50% inhibition of binding; Fmoc, 9-fluorenylmethoxycarbonyl; ELISA, enzyme-linked immunosorbent assay; t.u., transducing unit.

clones from the decapeptide library. K91 infected with library phage were plated on a total of 10 plates (150 mm × 15 mm) of LB medium containing 20 µg tetracycline/ml. The colonies were transferred and cross-linked to nylon membranes under u.v. light (Stratagene Duralon-UV). Treatment of the membranes and u.v. cross-linking were performed as described by the manufacturer (Stratagene, UV Stratalinker 1800; 1200 µJ, 30 s). The membranes were pre-hybridized at 42°C in 5 × SSC (750 mM-NaCl, 75 mM-trisodium citrate, 40 mM-sodium phosphate (pH 7.5), hybridization buffer (5 × Denhardt's solution; albumin/polyvinylpyrrolidone/Ficoll each at 1 mg/ml diluted in 5 × SSC containing 100 µg denatured salmon sperm DNA/ml). Oligonucleotide probes were phosphorylated as described by Sambrook *et al.* (1989). Following pre-hybridization, labeled probe (2.0 × 10⁶ cts/min per ml) was added to the pre-hybridization solution and the hybridization allowed to proceed overnight at 42°C with gentle agitation. The filters were washed 3 times in 0.1% (w/v) SDS, 2 × SSC at 42°C with gentle agitation (10 min wash). The filters were rinsed briefly in water and then in washing buffer: 3 M-tetramethylammonium chloride (TMAC), 50 mM-Tris·HCl (pH 8.0). The filters were washed in a shaking water-bath in washing buffer (1 h washes). The washing temperature that selected for perfect hybrids was determined empirically by use of a set of control filters. Each control set consisted of filter-lifts of colonies containing control phage with appropriate inserts: one insert was a perfect match to the oligonucleotide probe, and the other differed in sequence by a single base from a perfect match to the oligonucleotide probe. After washing, the filters were placed under film (Kodak XAR-2) for 4 h. After analysis of the control filters, the washing step was repeated, and the washing temperature increased sequentially in 2 deg.C increments from 50°C, until a temperature was reached at which all but perfect matches were eliminated, as revealed by the control filters. Washing temperatures approaching 56°C selected for perfect hybrids for all 4 probes, as was predicted by Wood *et al.* (1985). Positive clones from the decapeptide library that hybridized to the probes were confirmed as perfect matches by DNA sequencing.

(d) Affinity selection

The decapeptide library was subjected to a modification of the affinity selection technique of Parmley & Smith (1988). For comparison, a hexapeptide library donated by G. Smith was affinity-selected. The monoclonal antibody used for affinity selection was generated by immunization with recombinant Env 2-3 of human immunodeficiency virus (SF-2 isolate of HIV-1; Chapman *et al.*, 1991). The monoclonal antibody (mAb) was mapped to a 10 residue epitope (RAFHTTGRII) on the v3 loop of gp120 (Haigwood *et al.*, 1990). The antibody was biotinylated with biotin-XX-NHS (Calbiochem) as described by Parmley & Smith (1988).

Two rounds of affinity selection were performed. Streptavidin-coated, oxirane-derivatized polyacrylamide beads (Bass *et al.*, 1990) were used for the first round of affinity selection to allow an excess of streptavidin sites (and therefore the number of antibody sites) over the number of phage particles. The beads (Sigma, 1 µm particle size) were coated with streptavidin by incubating 500 mg of beads with 1 mg streptavidin/ml in 1 ml of phosphate-buffered saline (PBS) for 48 h at room temperature with gentle agitation. The beads were blocked with a solution of 1 mg bovine serum albumin/ml

for 24 h at 4°C, followed by 10 ml of 5% (v/v) β-mercaptoethanol in PBS for 24 h at 4°C with gentle agitation. The beads were then washed 5 times with 10 ml of PBS and stored in PBS containing 1 mg bovine serum albumin/ml.

The first round of affinity selection was performed by incubating 50 µl of biotinylated IgG (7.5 µg) with the streptavidin-coated beads (10 mg, 10 nmol/g resin) for 1 h at room temperature with agitation (a final antibody concentration of 1.0 µM). The beads were washed with PBS, 0.1% albumin. The library phage (10¹⁰ to 10¹¹ transducing units in 50 µl) were then added to the antibody-complexed beads and incubated for 2 h at room temperature with gentle agitation. The beads were washed by alternately suspending and pelleting 7 times with 400 µl of PBS, 0.5% (v/v) Tween 20. The bound phage were eluted with 0.4 ml of 0.1 M-glycine (pH 2.2), 1 mg albumin/ml. After elution, the pH of the phage suspension was adjusted to pH 7.0 with approx. 60 µl of 1 M-Tris·HCl (pH 9.0). The eluted phage were amplified after the first round of affinity selection, as described by Parmley & Smith (1988).

The second round of selection was done as described by Parmley & Smith (1988). This second round was performed under more stringent conditions, using fewer streptavidin sites. Polystyrene Petri dishes (60 mm × 15 mm) were coated with streptavidin (2.5 ml, 3 µg/ml in 0.1 M-NaHCO₃, pH 8.6) overnight at 4°C. The plates were then blocked with PBS, 1% albumin for 1 h at room temperature and washed with PBS (3 × 8 ml) just prior to use. The amplified phage library obtained from the first round of affinity selection (48 µl, 10⁹ to 10¹⁰ transducing units) and biotinylated antibody (20 µl, 750 ng, 250 nM) were incubated together for 2 h at room temperature (a final antibody concentration of 10 nM). The antibody/phage mixture was then diluted with 0.5 ml of PBS and added to the streptavidin-coated plates. The solution was incubated for 30 min at room temperature with gentle agitation. The plates were washed 8 times at 5 min intervals with PBS/0.5% Tween 20 (7 ml/wash). The bound phage were eluted and the solution neutralized as above.

(e) Immunological screening

The affinity-selected hexapeptide and decapeptide libraries were screened by an immunological colony screening technique adapted from Sambrook *et al.* (1989). Clones from each round of affinity selection were plated on 15 mm × 150 mm LB medium with 20 µl tetracycline/ml plates, at a density of 500 to 1000 well-separated colonies/plate, and were transferred to nitrocellulose membrane filters. The filters were immediately washed twice in TNT buffer (10 mM-Tris·HCl (pH 8.0), 150 mM-NaCl, 0.05% Tween 20 with no lysis step (30 min wash). The filters were blocked with blocking buffer (20% normal goat serum diluted in TNT buffer) for 30 min at room temperature with gentle agitation on a rotary platform. The anti-gp120 antibody was diluted 1:1000 (v/v) in blocking buffer to a final volume of 15 ml/filter and incubated with the filters for 2 h at room temperature with gentle agitation. The filters were then washed sequentially in washing buffer A (TNT buffer containing 0.1% albumin), washing buffer B (TNT buffer containing 0.1% albumin, 0.1% Nonidet P-40) and washing buffer A for 10 min at room temperature with gentle agitation. The secondary antibody, goat anti-mouse IgG-horseradish peroxidase conjugate (Boehringer-Mannheim), was diluted 1:200 (v/v) in blocking buffer to a final volume of 15 ml/filter and incubated with the filters for 2 h at room temperature

Table 1
Control bacteriophage for binding anti-gp120

	Insert sequence†	IC50‡
T bacteriophage	RAFHTTGRII	6.3 nM
I bacteriophage	RAFHTTGAII	320 nM
L bacteriophage	FHTTGRII	3.9 µM

† Amino acid sequence of the insertions present in the pIII protein of the control bacteriophage.

‡ The IC50s were measured by competition ELISA of the free peptide insert, as described in the text.

with gentle agitation. The filters were then washed as above. Positive clones were located by color development using 3,3'-diaminobenzidine tetrahydrochloride, as described by the manufacturer (Pierce). Prior to sequencing, positive clones were verified by a second round of immunological screening. Three control bacteriophage that bind to the anti-gp120 monoclonal antibody with a range of IC50s between 4 µM and 6 nM were screened simultaneously with the libraries as positive controls (Table 1).

(f) Analysis of candidate peptide ligands

Candidate peptides (*N*-acetylated, *C*-amidated) were synthesized on an automated synthesizer using Fmoc chemistry (Zuckermann *et al.*, 1992). The peptides were assayed in a competition ELISA format over a concentration range of 0.1 nM to 500 µM. Immulon 1 microtiter plates (Dynatech Laboratories) were coated overnight at 4°C with recombinant gp120 antigen at a concentration of 0.2 µg/well in 50 mM-sodium borate (pH 9.0). Then 50 µl of each peptide was added to the wells, followed by the addition of 50 µl of diluted mAb (26 µM-stock diluted 1:50,000 (v/v)). The mixtures were incubated in 0.5 M-NaCl, 1% (v/v) Triton X-100, 0.1% (w/v) casein, 250 mM-sodium phosphate (pH 7.5) for 1 h at 37°C. The plates were washed 6 times with washing buffer (150 mM-NaCl, 0.5% Triton X-100) and incubated with 100 µl of horseradish peroxidase-conjugated goat anti-mouse IgG antibody (1 mg/ml stock diluted 1:1000 (v/v), Boehringer-Mannheim) for 1 h at 37°C. The plates were washed as above and the bound conjugated antibody was quantified by color development with 100 µl of *o*-phenylenediamine (5 mg/ml in 50 mM-sodium citrate, 0.02% (v/v) H₂O₂, pH 5.1). Plates were read at 450 nm on a Molecular Devices ThermoMax microplate reader.

3. Results

(a) Construction of a bacteriophage decapeptide library

The decapeptide library was constructed using a self-priming oligonucleotide, which increases the efficiency of both the annealing and priming reactions. Transformation of the library into DH10B cells was accomplished by electroporation. Electroporation of the decapeptide library yielded 4×10^8 independent transformants, at an efficiency of about 2×10^7 transformants/µg. The fraction of clones containing a library insert was determined by comparing the ligation frequency to the religation frequency. The frequency of vector religation was 10^2 -fold lower than the ligation frequency. Following amplification, the library was found to

5'GGCCGACGTGGCC(NNG/T)₁₀GCGGCCTCTGGGGCC 3'

Probe 3'A AATCGTGCGGCCTCTGG 3'

Probe 3'B IATCGTGC GGCCCTCTGG 3'

Probe 5'A CCGACGTGGCCTATCGG3'

Probe 5'B CCGACGTGGCCTATCGI 3'

Figure 2. The DNA sequences of the 4 oligonucleotide (17 bp) probes used to evaluate the diversity of the decapeptide library. Each of the probes contains 11 bp complementary to either the 5' or 3' constant, flanking sequence of the insert and 6 bp (NNG/T NNG/T; bold face) complementary to the corresponding 5' or 3' end of the 30 bp variable library sequence. Probes 5'A and 5'B differ by a single base-pair of the variable library region, as do probes 3'A and 3'B (underlined).

contain about 6×10^{12} transducing units (t.u.), or approximately 1.5×10^4 copies of each possible bacteriophage.

(b) Genetic diversity of the decapeptide library

In order to evaluate the genetic diversity of the decapeptide library, a modified colony hybridization protocol utilizing tetramethylammonium chloride (TMAC) was developed that allows the discrimination of full-length and partial hybrids (i.e. perfect *versus* imperfect matches).

The sequences of the four 17 bp oligonucleotide probes used to evaluate the diversity of the decapeptide library are shown in Figure 2. Each of the probes contains 11 bp complementary to either the 5' or 3' constant, flanking sequence of the insert and 6 bp (NNG/T NNG/T) complementary to the corresponding 5' or 3' end of the 30 bp variable library sequence. Probes 5'A and 5'B differ by a single base-pair in the variable library region, as do probes 3'A and 3'B. Based on the frequency of each possible base-pair at each of the six positions of the sequence (NNG/T NNG/T) of the degenerate region of the probe, the expected number of perfect matches out of the 5000 colonies from the decapeptide library for each of the four probes 3'A, 3'B, 5'A and 5'B is approximately five. The observed number of positives for each probe roughly correlated with the theorized number: 5'A, four positive clones; 5'B, three positive clones; 3'A, three positive clones; and 3'B, four positive clones. As expected, the four probes identified discrete positive clones from the same filter lifts. The three clones that hybridized to the 3'A probe were confirmed as perfect matches by DNA sequencing.

(c) Affinity selection

The yield of bacteriophage, defined as the number of bacteriophage obtained in an elution fraction relative to the initial number of bacteriophage, was determined for each round of affinity selection performed on the hexapeptide and decapeptide libraries (Table 2). A significant increase (1000-fold)

Table 2

The yield of phage following two rounds of affinity selection of the hexapeptide and decapeptide libraries

Library	Selection round	Phage yield (%)
Hexapeptide	1	1.2×10^{-4}
	2	2.2×10^{-1}
Decapeptide	1	3.7×10^{-4}
	2	9.5×10^{-1}

in the yield of bacteriophage from the first round of affinity screening *versus* the second round was observed for both libraries (Table 2). This increase in yield suggests an enrichment of mAb-binding ligands. Random and immunological screenings were performed on the eluted bacteriophage to further confirm this enrichment.

(d) *Random screening versus immunological screening*

An increase in the percentage of positive clones between the first and second rounds of selection of both the hexapeptide and decapeptide libraries was observed using immunological screening, as expected from successful affinity selection (see Table 5).

Clones from the second round of affinity selection of the hexapeptide library were randomly chosen for sequencing. Of the 120 clones sequenced, there were 50 distinct peptide sequences (data not shown). Twelve of the peptide sequences were represented more than once. Two of the 12 peptide sequences were represented by clones with differing nucleotide sequences, strongly suggesting that they are not sibling clones. These 12 sequences and eight unique sequences (Fig. 3) were synthesized as 14mer peptides, each containing the hexapeptide library insert sequence and the eight flanking amino acids from the bacteriophage pIII protein




Figure 3. Twenty of the 50 hexapeptide sequences obtained from the random screening of 120 clones, grouped by the frequency with which each appeared. Twelve separate peptide sequences are represented more than once. The 20 peptides were synthesized and assayed for binding with anti-gp120 mAb. Of the 20 free peptides, 10 have an IC₅₀ of >10 μM (hatched box) and 10 have an IC₅₀ <10 μM (filled box).

Table 3

IC₅₀s of hexapeptide library epitopes for binding anti-gp120

Insert sequence†	IC ₅₀ ‡	Frequency per 120 clones
ESTRPM	500 nM	3
PCCRAF	630 nM	7
ESTRAM	1.3 μM	1
ESTRRM	1.6 μM	8§
KTFYCC	2.5 μM	3
SCCRPM	2.5 μM	1
CCRGGP	2.5 μM	1
PCCRAP	4.0 μM	1
GETRRM	5.0 μM	9§
DCCRTF	15 μM	1

† Amino acid sequence of the hexapeptide insertions present in the pIII protein of the library bacteriophage.

‡ The IC₅₀s of the free peptides were measured by competition ELISA, as described in the text.

§ The DNA sequences of these isolates differ.

(ADGA[X]₆GAAG). The IC₅₀s of the peptides were measured by competition ELISA. Ten of the 20 peptides competed with antigen for binding to anti-gp120 mAb with IC₅₀s of less than 10 μM (Table 3). Two common subsequence motifs were observed: XSTRXM and XCCRX. These motifs appeared in 29% of the 120 clones.

Immunological screening was performed on the hexapeptide library after the second round of affinity selection. Of the 27 immunopositive clones that were chosen for sequencing, 25 contained sequences identical with the submicromolar (IC₅₀) binders identified by random screening (Table 4). Therefore,

Table 4

IC₅₀s of library epitopes chosen by immunoscreening for binding anti-gp120 mAb

Library	Insert sequence†	IC ₅₀ ‡	Frequency
Hexapeptide	ESTRPM	500 nM	1
	PCCRAF	630 nM	16
	ESTRRM	1.6 μM	7
	SCCRPM	2.5 μM	1
	VSTRPM	N/A	1
	GSTRRM	N/A	1
Decapeptide	ESTRPMAGDG	300 nM	1
	RALESTRAME	100 nM	1
	PSTRPMQRSG	>10 μM	1
	EIRASTRPMR	500 nM	1
	QRSLCCREFF	100 nM	1
	FCCRMPSSGA	1 μM	1
	ECCRPMPHRH	1 μM	1
	SSRPMRLTKT	300 nM	1
	PCKDLSRF	3 μM	1
	STGPMRPLQV	600 nM	1
	SSDRAWWCCS	300 nM	1
	GRAWGVARDR	1 μM	1
	LARKDGFGGW	300 nM	1
	SIVSVREVF	>10 μM	1

† Amino acid sequence of the hexamer and decamer insertions present in the pIII protein of the library bacteriophage selected by *in situ* immunoscreening. Bold face sequences are either XSTRXM or XCCRX, or are the sequences of known binders.

‡ The IC₅₀s of the free peptides were measured by competition ELISA, as described in the text. N/A, not assayed.

Table 5
Immunopositive clones per round of affinity selection of libraries

Library	Round	Frequency (%)
Hexapeptide	1	2.6
	2	17
Decapeptide	1 pre-amplification	7.2
	1	8.8
	2 pre-amplification	22
	2	25

at least 93% of the clones selected by immunological screening of the hexapeptide library contained peptides with IC₅₀s less than 10 μ M, and all either had the XCCR_X or XSTR_{XM} motifs as compared with random screening, which yielded a low proportion of binding peptides, 29%. This lower rate could have been predicted by immunological screening, which indicated a 17% frequency of immunopositive clones (Table 5).

Similar results were obtained from the immunological screening performed on the decapeptide library (Table 5). Four of the 14 immunopositive clones selected for sequencing had the XSTR_{XM} motif, and three had the XCCR_X motif (Table 4). The remainder of the clones contained unique novel sequences. All 14 sequences were synthesized as free decapeptides and the IC₅₀s of the peptides were measured by competition ELISA.

In order to determine the limit of detection of immunological screening protocol, the three positive control bacteriophage (T, I and L, Table 1) were screened. Only the submicromolar (IC₅₀) bacteriophage-infected colonies (T and I) were detectable with the protocol described above. Therefore, the limit of detectability for the immunological screening technique with the anti-gp120 antibody is an IC₅₀ of approximately 4 μ M. Only the T bacteriophage-infected colonies were detected when the concentration of the anti-gp120 antibody was decreased tenfold (1:10,000 (v/v) dilution). Therefore, it should be possible to differentiate between binders with IC₅₀s of 1 μ M and 10 nM, by appropriately lowering the concentration of the primary antibody used in the immunological screening.

4. Discussion

We have developed a strategy for generating peptide libraries of great size and complexity utilizing a self complementary, self-priming oligonucleotide. The simplicity of this technique permits the rapid, efficient construction of phage libraries. The decapeptide library constructed using this strategy was large enough to ensure that a complete set of hexamer peptide sequences were represented in the five possible hexamer frames within the decapeptide sequence (a phage library of 4×10^8 with 5 hexamer frames encompasses greater than 1×10^9 hexamers, or $(32)^6$), allowing comparison of affinity-selected peptide sequences from a complete hexamer

library versus the smaller hexapeptide library (donated by G. Smith). Half of the affinity-selected immunopositive decapeptide library phage chosen for sequencing contained the same two hexamer motifs found in the hexapeptide library, indicating that the anti-gp120 antibody recognized the linear hexamer sequence in the decapeptide library context. In addition, free hexapeptides containing these sequences had equivalent IC₅₀s to those of the free decapeptides (data not shown), indicating that the flanking residues have no effect on the binding of these hexamers.

Neither of the hexamer motifs XSTR_{XM} nor XCCR_X is related to the sequence of the original gp120 epitope sequence RAFHTTGR_{II}, and neither these peptide ligands nor the unrelated peptide ligands selected by immunological screening (Table 4) had a equivalent or higher IC₅₀ than the original epitope. It is possible that we are reproducing the same phenomenon seen by others (Scott & Smith, 1990; Cwirla *et al.*, 1990); namely, that peptides of high affinity (nM IC₅₀s) have no selective advantage over peptides of moderate affinity (μ M IC₅₀s) under the conditions used to date. It is probable that the epitope RAFHTTGR_{II} was not discovered because the decapeptide library contains only 4×10^{-5} % of a complete decapeptide library.

Like others, we have identified epitopes unrelated in sequence to the known epitope of a monoclonal antibody. Scott & Smith (1990) isolated the epitope CRFVWC by affinity selection with the monoclonal antibody M33, although it bears no sequence similarity to the eliciting myohemerythrin epitope DFLEKI. It can therefore be conjectured that an unrelated combination of residues might approximate the binding valencies and spatial orientation of the residues particular to an eliciting epitope. That unrelated peptide sequences can be affinity-selected by a monoclonal antibody would also imply that the use of peptide libraries for epitope mapping might not be as unequivocal as previously thought. However, it also suggests that phage peptide libraries can be used to great advantage in other applications, such as to investigate cross-reactivities of antibodies.

The presence of a genetic bias in a library decreases the effective size of the library, thereby decreasing the probability that a specific clone will be isolated during screening. Genetic bias can be introduced at many steps during library construction; for example, during synthesis of the variable positions within the library insert, during cloning or as a result of biological selection during library amplification. It is therefore preferable to examine the genetic diversity of a bacteriophage library following its final amplification.

The genetic diversity of the decapeptide library was analyzed using a modified colony hybridization technique (Wood *et al.*, 1985). TMAC mediates a shift of the dissociation temperature (t_d) of A·T base-pairs to that of G·C base-pairs, with the result that the binding of an oligonucleotide probe becomes solely dependent on the length of the

hybrid. Therefore the use of TMAC allows the discrimination of full-length and partial hybrids (i.e., perfect *versus* imperfect matches).

Two sets of oligonucleotide probes were used to examine the genetic diversity of the decapeptide library. These sets were good controls for identifying perfect matches, as the sequences differed only at the final base. Mismatches at the ends of hybrids have less effect on hybrid stability than internal mismatches, therefore, the ability of this method to select for perfect matches was demonstrated unequivocally.

Of the two sets of probes, one set was designed to scan the 5' and the other the 3' termini of the variable region in the library insert. Oligonucleotides are synthesized in a 3' to 5' direction. Any bias in the sequence of the degenerate region of the library insert due to non-random nucleotide incorporation would be most obvious at the 5' end of the oligonucleotide, as the effect of an inefficient coupling reaction would be amplified. The analysis of diversity at the 3' terminus, which is unlikely to show bias, served as a control for the measurements of bias at the 5' terminus.

If bias introduced during oligonucleotide synthesis was due to the cumulative effect of the inefficient coupling of a single base at every coupling of that base, this bias would be most easily observed if the probes were multimers of single bases (for example, a poly(dG) probe). For reasons of convenience, the probes used in examining the diversity of the decapeptide library were not multimers of single nucleotides. Therefore, the analysis of bias of the decapeptide library was of only moderate stringency. However, this technique can yield a quantitative analysis of the genetic bias of a library if monobase multimers are used as probes.

Affinity selection enriches for the peptide ligands present in a pool of random peptides but it does not eliminate all non-specific peptides, such as the peptides that bind to streptavidin (Devlin *et al.*, 1990). Of the 120 clones selected by the random screening of the hexapeptide library 30% contained the hexamer sequence WRANGT (Fig. 3). Although affinity selection enriched for this hexamer, the free decapeptide containing this sequence did not compete with antigen for binding to anti-gp120 mAb. Interestingly, WRANGT was not isolated during the immunological screening of either the hexapeptide or decapeptide libraries, indicating the effectiveness of the immunological screening technique at identifying binding-site associated phage.

The standard method for screening the output of an affinity selection experiment is the sequencing of random clones followed by analysis of the peptide sequences for motifs prior to further investigation. Felici *et al.* (1991) have described the limited (48 clones) confirmatory immunological testing of affinity-selected bacteriophage *via* dot blot. However, a more efficient approach is to rapidly screen large quantities of clones ($>10^4$) after affinity selection by using a colony immunological screening technique that highlights binding clones. A smaller

number of clones therefore needs to be sequenced, since these candidates are likely to bind with a micromolar IC50. At least 93% of the clones isolated by immunological screening of the hexapeptide library contained peptide sequences that bind with IC50s $<10 \mu\text{M}$. In comparison, random screening yielded a low percentage of binding peptides, 29%. Of the 14 immunopositive clones selected from the decapeptide library, 86% of the identified peptides competed with antigen for binding to anti-gp120 mAb with IC50s $<10 \mu\text{M}$.

In conclusion, we have generated a peptide library of great size and complexity using a simple and efficient self-priming oligonucleotide scheme. We have examined its genetic diversity by using a modified colony hybridization protocol and subjected it to affinity selection with an anti-gp120 monoclonal antibody. The immunological screening technique described in this publication represents an improvement in bacteriophage library screening as it allows the rapid, direct identification of binding bacteriophage. The immunological screening of affinity-selected libraries is therefore a powerful tool for choosing appropriate peptide ligands as candidates for further investigation.

We are grateful to Dr G. Smith and Dr J. Scott for providing the hexapeptide bacteriophage library and Dr N. Haigwood for providing the anti-gp120 antibody. We thank Drs F. Cohen and S. Rosenberg for helpful discussion, and H. Lukatch and S. Banville for technical assistance.

References

- Bass, S., Greene, R. & Wells, J. A. (1990). Hormone phage: an enrichment method for variant proteins with altered binding properties. *Proteins*, **8**, 309-314.
- Broome, S. & Gilbert, W. (1978). Immunological screening method to detect specific translation products. *Proc. Nat. Acad. Sci., U.S.A.* **75**, 2746-2749.
- Chapman, B. S., Thayer, R. M., Vincent, K. A. & Haigwood, N. L. (1991). Effect of intron A from human cytomegalovirus (Towne) immediate-early gene on heterologous gene expression in mammalian cells. *Nucl. Acids Res.* **19**, 3979-3986.
- Clackson, T., Hoogenboom, H. R., Griffiths, A. D. & Winter, G. (1991). Making antibody fragments using phage display libraries. *Nature (London)*, **352**, 624-628.
- Cwiria, S. E., Peters, E. A., Barrett, R. W. & Dower, W. J. (1990). Peptides on phage: a vast library of peptides for identifying ligands. *Proc. Nat. Acad. Sci., U.S.A.* **87**, 6378-6382.
- Devlin, J. J., Panganiban, L. C. & Devlin, P. E. (1990). Random peptide libraries: a source of specific protein binding molecules. *Science*, **249**, 404-406.
- Dower, W. J., Miller, J. F. & Regsdale, C. W. (1988). High efficiency transformation of *E. coli* by high voltage of electroporation. *Nucl. Acids Res.* **16**, 6127-6145.
- Felici, F., Castagnoli, L., Musacchio, A., Jappelli, R. & Cesareni, G. (1991). Selection of antibody ligands from a large library of oligopeptides expressed on a multivalent exposition vector. *J. Mol. Biol.* **222**, 301-310.

- Güssow, D. & Clackson, T. (1989). Direct clone characterization from plaques and colonies by the polymerase chain reaction. *Nucl. Acids Res.* **17**, 4000.
- Haigwood, N. L., Shuster, J. R., Moore, G. K., Lee, H., Skiles, P. V., Higgins, K. W., Barr, P. J., George-Nascimento, C. & Steimer, K. S. (1990). Importance of hypervariable regions of HIV-1 gp120 in the generation of virus neutralizing antibodies. *AIDS Res. Hum. Retrovir.* **6**, 855-869.
- Lowman, H. B., Bass, S. H., Simpson, N. & Wells, J. A. (1991). Selecting high-affinity binding proteins by monovalent phage display. *Biochemistry*, **30**, 10832-10838.
- Maniatis, T., Kee, S. G., Efstratiadis, A. & Kafatos, F. C. (1976). Amplification and characterization of a β -globin gene synthesized *in vitro*. *Cell*, **8**, 163-168.
- Marks, J. D., Hoogenboom, H. R., Bonnert, T. P., McCafferty, J., Griffiths, A. D. & Winter, G. (1991). By-passing immunization: human antibodies from V-gene libraries displayed on phage. *J. Mol. Biol.* **222**, 581-597.
- Miller, J. H. (1972). *Experiments in Molecular Genetics*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Parmley, S. F. & Smith, G. P. (1988). Antibody-selectable filamentous fd phage vectors: affinity purification of target genes. *Gene*, **73**, 305-318.
- Sambrook, J., Fritsch, E. F. & Maniatis, T. (1989). *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Sanger, F. S., Nicklen, S. & Coulson, A. R. (1977). DNA sequencing: with chain-terminating inhibitors. *Proc. Nat. Acad. Sci., U.S.A.* **74**, 5463-5467.
- Scott, J. K. & Smith, G. P. (1990). Searching for peptide ligands with an epitope library. *Science*, **249**, 386-390.
- Wood, W. I., Gitschier, J., Lasky, L. A. & Lawn, R. M. (1985). Base composition-independent hybridization in tetramethylammonium chloride: a method of oligonucleotide screening of highly complex gene libraries. *Proc. Nat. Acad. Sci., U.S.A.* **82**, 1585-1588.
- Young, R. A. & Davis, R. W. (1983). Efficient isolation of genes by using antibody probes. *Proc. Nat. Acad. Sci., U.S.A.* **80**, 1194-1198.
- Zuckermann, R. N., Kerr, J. M., Siani, M. A. & Banville, S. C. (1992). Design, construction and application of a fully automated equimolar peptide mixture synthesizer. *Int. J. Pept. Protein Res.* in the press.

Edited by J. Karn